

Berry's Phase, Josephson's Equation, and the Dynamics of Weak Link Superconductors and Their Vortices

Frank Gaitan and Subodh R. Shenoy

International Center for Theoretical Physics; P. O. Box 586; Miramare; 34100 Trieste, ITALY
(February 1, 2008)

We examine the dynamical consequences of Berry's phase for Josephson junctions, junction arrays, and their vortices. Josephson's equation and the related phase slip voltages are shown to be unaffected by Berry's phase. In an annular Josephson junction, Berry's phase is seen to generate a new current drive on a vortex. In the continuum limit, vortex dynamics in a 2D array is shown to map onto that of a 2D film. A Hall sign anomaly is expected in arrays; and the merits of arrays for studies of disorder on vortex motion is discussed.

74.50.+r, 74.20.-z, 03.65.Bz

In a very interesting paper [1], Ao and Thouless showed that vortex motion in a Type-II superconductor generates a Berry phase in the many-body ground state, and as a consequence, the Magnus force of classical hydrodynamics acts on the vortex [2]. Their work led others to calculate the effective action for the superconducting condensate S_g using the BCS model [3,4]. It was shown by one of these authors [4] that when the Berry phase is non-removable, it manifests as a Wess-Zumino (WZ) term in S_g . Ref. [4] also pointed out that a WZ term might arise in the effective action of a Josephson Junction (JJ) when the superconducting gap phase ϕ contained a moving vortex singularity.

In this Letter we examine the consequences of the Berry-phase-induced WZ term for the dynamics of weak link superconductors and their vortices. As far as we are aware, this is the first time such a study has been carried out, either for single weak link systems, or for an array of such links [5]. We work within the confines of the BCS model of superconductivity; assume the gap phase contains a moving vortex singularity $\phi(\mathbf{r} - \mathbf{r}_0(t))$, where $\mathbf{r}_0(t)$ is the vortex trajectory; and $T = 0$. A detailed presentation of this work will be given elsewhere [6]. The essential results of this paper are: (1) in a single superconducting grain, the Josephson-Gorkov equation is not modified in any physically relevant way by the WZ term, and consequently, phase slip voltages are not effected by Berry's phase; (2) Berry's phase leads to a new current drive acting on a vortex in an annular JJ, though it proves ineffective (in this type of junction) for geometrical reasons; (3) in a 2D Josephson Junction Array (JJA), this new Berry-phase-induced current drive *is* operative; also, in the Continuum Limit (CL), the superconducting (SC) dynamics of the JJA maps onto that of a 2D film so that a Hall Effect sign anomaly is expected in a JJA if the grains are made of a material which shows such an anomaly in a 2D film; and (4) a JJA in the CL appears ideally suited to test recent predictions of Vinokur et. al. [7] regarding the Hall conductivity and resistivity.

(i) Single Superconducting Grain: We begin by considering a single superconducting grain in which a moving vortex singularity is present. The effective action for the superconducting condensate S_g was obtained in Ref. [3], though our notation will follow Ref. [4]. This action contains space-time gradients up to second order and has the form $S_g = S_0 + S_{wz} + S_2$. S_0 corresponds to the action for a uniform gap function and depends only on $|\Delta|$, whereas S_{wz} and S_2 depend *only* on the gap phase $\phi = \phi(\mathbf{r} - \mathbf{r}_0(t))$. The gap equation corresponds to the Euler-Lagrange equation (EL-eqn) obtained by variation of S_g (viz. S_0) with respect to $|\Delta|$ [8]. This equation acts to fix $|\Delta|$. Outside the critical region near T_c , fluctuations away from the extremal solution of the gap equation can be ignored, so one can treat $|\Delta|$ as fixed, and approaching Δ_0 far from the vortex core. Beyond fixing $|\Delta|$, S_0 will not interest us and so will not be written out explicitly. The remaining contributions to S_g are [3,4]

$$S_{wz} + S_2 = \int d^2x dt \left[\rho_s \left(\frac{\hbar}{2} \partial_t \phi + eA_0 \right) + N(0) \left(\frac{\hbar}{2} \partial_t \phi + eA_0 \right)^2 + \frac{m\rho_s}{2} \mathbf{v}_s^2 + \frac{1}{8\pi} \left\{ (\mathbf{H} - \mathbf{H}_{ext})^2 - \mathbf{E}^2 \right\} \right] . \quad (1)$$

S_{wz} is the WZ term and corresponds to the term in eqn. (1) whose integrand is gauge invariant and first order in time derivatives of the gap phase ϕ . S_2 is the remaining collection of terms. Here ρ_s is the density of superconducting electrons; A_0 is the scalar potential induced by the vortex motion; $N(0)$ is the normal density of states at the Fermi surface for one spin component; m is the electron mass; $\mathbf{v}_s = \hbar(\nabla\phi - (2e\mathbf{A}/\hbar c))/2m$ is the superfluid velocity; $\mathbf{H} = \nabla \times \mathbf{A}$ is the microscopic magnetic field; $\mathbf{H}_{ext} = \nabla \times \mathbf{A}_{ext}$ is the externally applied magnetic field; $\mathbf{E} = -\nabla\mathcal{E} = -\nabla A_0 - \partial(\mathbf{A}/c)/\partial t$ is the electric field generated by the vortex motion; and $\mathcal{E} = -\int^\mathbf{x} d\mathbf{l} \cdot \mathbf{E}$ is the gauge invariant electromotive force whose gradient determines $\mathbf{E}$.

If we momentarily consider the case of a superconducting grain in which no vortex is present, no WZ term will appear in eqn. (1) [4]. This case was considered in Ref. [8] and the Josephson-Gorkov equation (JG-eqn) was obtained

by noting that the remaining term containing $\partial_t \phi$ is positive definite and thus contributes least to S_g when it is zero. It follows from this remark that $-2eA_0 = \hbar \partial_t \phi$ (viz. the JG-eqn). Returning to the case of a grain containing a moving vortex, here the WZ term is present. One can combine the integrands of the first two terms in eqn. (1) to obtain $\{-\rho_s^2/4N(0)\} + N(0)\{\hbar \partial_t(\phi/2) + eA_0 + (\rho_s/2N(0))\}^2$. If one were to naively carry over the argument of Ref. [8], one would argue that this expression was smallest when the positive definite second term vanished, and subsequently, the “JG-eqn” (in the presence of a moving vortex) would be $-2eA_0 = \hbar \partial_t \phi(\mathbf{r} - \mathbf{r}_0(t)) + \rho_s(\mathbf{r})/N(0)$. On the basis of this analysis one would conclude that, for a given vortex trajectory $\mathbf{r}_0(t)$, the scalar potential A_0 was modified as a result of the Berry-phase-induced WZ term. Because of the intimate connection between the JG-eqn and the voltage differences induced by vortex motion through phase slip, such a modification, if true, might be thought to lead to new physics. In fact, the above argument is not correct for the case of a grain containing a moving vortex singularity, as we now show. The essential point is that, when a moving vortex is present, $\mathbf{E} \neq 0$. To obtain the JG-eqn, we vary S_g with respect to $\mathcal{E}$ (recall $\mathbf{E} = -\nabla \mathcal{E}$) to obtain Poisson’s equation

$$-\nabla^2 \mathcal{E} = 4\pi \left[2N(0)e \left\{ \frac{\hbar}{2} \partial_t \phi + eA_0 + \frac{\rho_s}{2N(0)} \right\} \right] . \quad (2)$$

The term in square brackets on the RHS of this equation must equal the electric charge density $e(\rho_s - \rho_0)$, where ρ_0 is the smeared out charge density of the positive ions in the lattice. One can obtain a consistency check on this identification by obtaining the EL-eqn for ϕ which is the continuity equation for electric charge. One finds [6] that the quantity in square brackets in eqn. (2) again appears as the charge density. Carrying out this identification yields the JG-eqn (in the presence of a moving vortex), $-2eA_0 = \hbar \partial_t \phi + \rho_0/N(0)$. Thus the JG-eqn does receive a correction term, although this correction is a physically irrelevant constant, and will not effect voltage differences. It can safely be discarded since the scalar potential is only defined modulo a constant. Note that for the case considered by Ref. [8], no WZ term occurs, and $\nabla \mathcal{E} = 0$; thus Poisson’s equation will not contain the final term in the curly brackets on the RHS of eqn. (2), and the LHS will vanish. Thus, in this case, we recover the usual JG-eqn from our approach. We also see that the argument of Ref. [8] (appropriate for the situation considered there) is not generally applicable, being inconsistent with Poisson’s equation when a moving vortex is present. Thus we obtain our first result: Berry’s phase does *not* modify the Josephson-Gorkov equation. Consequently, the usual arguments concerning phase slip voltages are also unaltered [9]. As first pointed out in Ref. [1], Berry’s phase *does* effect the motion of a vortex via the Magnus force. In Ref. [4], the derivation of this force from the WZ term is given, and it appears that this is the only effect of Berry’s phase in the case of a single superconducting grain.

(ii) *Single JJ*: We go on now to the case of a Josephson Junction in which two superconducting grains (L, R) are coupled through a weak link by the Josephson effect. We assume the weak link to be a tunneling barrier (TB). The Hamiltonian for this system is a sum of the individual grain Hamiltonians and two interaction terms which dynamically couple the grains: (i) a Coulomb term which accounts for the capacitive coupling of the electric charges across the TB; and (ii) a tunneling term responsible for coupling the gap phases of the grains [10]. The action for the JJ takes the form $S_{JJ} = S_g(L) + S_g(R) + S_c + S_t$ [8]. Here $S_g(i)$ ($i = L, R$) is the single grain action described above eqn. (1); and

$$S_c = - \int \frac{dt}{\hbar} \frac{C}{2} \left[\frac{\hbar}{2e} \partial_t \gamma \right]^2 ; \quad S_t = \int \frac{dt}{\hbar} \left[-I_c \frac{\hbar}{2e} \cos \gamma \right] .$$

C is the junction capacitance; $\gamma = (\phi_L - \phi_R) - \int_1^2 \mathbf{A} \cdot d\mathbf{l}$ is the gauge invariant (gap) phase difference across the TB; $I_c = \pi \Delta_0 / 2eR_n$ is the critical current, and R_n is the tunnel junction normal resistance. We restrict ourselves to Large JJ’s so that localized regions of magnetic flux are possible (vortices) in the TB, and the vortex motion in a JJ is one dimensional. The grains are assumed to be 2D (unit thickness in the z-direction), so that the JJ is embedded in R^2 . Consequently, the TB maps onto a (finite) 1D segment $P \subset R^2$ which will be referred to as the parameter space since $\mathbf{r}_0(t) \in P$ is the set of parameters appearing in the grain Hamiltonians, and whose time dependence generates the Berry phases. Since the presence of the WZ term in the grain action occurs only when the Berry phases are non-vanishing, we must examine whether this situation is possible in a JJ.

It is enough to consider the Berry phases in the single particle states since the many-body states will be constructed from them. We first consider a Traditional (Large Straight) JJ (TJJ) in which the parameter space P_T is a 1D segment whose endpoints correspond to physically distinct points. P_T has the topology of the unit interval $I = [0, 1]$. Thus, the only closed loops C possible in P_T will originate from an arbitrary point p_0 ; go out to a point p_1 via a segment C_1 ; and return to p_0 along $-C_1$. Note that each point on C_1 is passed through twice. This forces the Berry phase f_B to vanish,

$$f_B = - \oint_{C_1 - C_1} d\tau \left[i\dot{\mathbf{r}}_0 \cdot \langle E(\mathbf{r}_0) | \nabla_{\mathbf{r}_0} | E(\mathbf{r}_0) \rangle + \frac{e}{\hbar} A_0 \right] = 0 \quad .$$

This follows since $\langle E | \nabla_{\mathbf{r}_0} | E \rangle$ and A_0 are single-valued functions over P_T so that $\int_{C_1}$ is equal and opposite to $\int_{-C_1}$ for all C_1 and for all single particle states. Consequently, f_B vanishes for all the states in the many-body Fock space, and no WZ term is generated in $S_g(i)$ by the vortex motion (for a TJJ). From the point of view of Berry's phase, a more interesting JJ is the Annular JJ (AJJ) in which the TB has a ring-like topology [11]. Here the TB maps onto the unit circle in R^2 so that $P_A = S^1$. Berry phases for closed loops that wind around S^1 are non-vanishing (in the gauge $A_0 = 0$, the BCS groundstate Berry phase induced by a single traversal of S^1 is $-\pi N_s$, where N_s is the number of superconducting electrons). Thus, for an AJJ, the Berry phases are non-trivial, and the grain actions each contain a WZ term whose consequences we now explore.

We restrict ourselves to the case where an applied (uniform) transport current flows through the AJJ so that $\mathbf{v}_s = \mathbf{v}_T + \mathbf{v}_{\text{circ}}$. Here $\mathbf{v}_T$ is the superflow velocity of the transport current; and $\mathbf{v}_{\text{circ}}$ is that of the superflow circulating about the vortex. We now show that Berry's phase modifies the current drive acting on a vortex. This drive describes the coupling of the grain condensates to the vortex, and the coupling arises from the terms in $S_g(i)$ linear in $\mathbf{v}_{\text{circ}}$. One such term comes from the $m\rho_s \mathbf{v}_s^2/2$ term common to the grain action of all types of JJ's. It leads to the familiar Lorentz current drive whose contribution to S_t is $\int dt (-I\gamma/2e)$ [8]. In an AJJ, the WZ term also contributes to the current drive term in S_t . We will refer to this contribution as the Magnus drive since its origin (the WZ term) is the same as that of the Magnus force [1,4]. Let $\alpha \in S^1$ parameterize position along the TB (recall $P_A = S^1$); and let $\hat{\mathbf{n}}_L(\alpha)$ ($\hat{\mathbf{n}}_R(\alpha)$) be the unit outward normal to the face of the Left (Right) grain which interfaces with the TB. In all case we are familiar with, $\hat{\mathbf{n}}_L(\alpha) = -\hat{\mathbf{n}}_R(\alpha) = \hat{\mathbf{n}}(\alpha)$. By partial integration of these coupling terms, one can show [6] that the total current drive contribution to S_t is

$$S_{dr} = \int dt dz r d\alpha \gamma(\alpha) \left[-\frac{\rho_s \hbar}{2} \hat{\mathbf{n}}(\alpha) \cdot (\mathbf{v}_T - \dot{\mathbf{r}}_0) \right] \quad . \quad (3)$$

The $\mathbf{v}_T$ -term is the Lorentz drive, and the $\dot{\mathbf{r}}_0$ -term is the Magnus drive. In principle, the Magnus drive is non-zero; though, in practice, tunnel junctions are usually constructed such that $\hat{\mathbf{n}}(\alpha) \perp \dot{\mathbf{r}}_0$, so that the Magnus drive effectively vanishes in a smooth annular geometry. Thus Berry phase effects in large JJ's are rather subtle. These effects are seen to be extremely sensitive to the topology of the tunneling barrier, and even where they are expected to occur (AJJ), the usual tunneling barrier geometry conspires to nullify its effects!

(iii) Junction Arrays: Finally, we consider a JJA which corresponds to a lattice L (lattice constant a_0) whose sites are occupied by SC grains. Phase coherence of the gap function throughout the array is established via the Josephson effect. Let i index the lattice sites, and $\{n_i\}$ index the nearest neighbors of i . Vortices in a JJA live on the dual lattice $\mathcal{L}$ which is the multiply-connected region separating the grains, and which is constituted by the TB's (i, n_i) between neighboring grains. Accordingly, the dual lattice is identified with the parameter space, $P = \mathcal{L}$. With respect to grain i , any closed loop C_i in $\mathcal{L}$ which winds around i will produce a non-vanishing Berry phase in this grain's SC groundstate. This will be true for all grains i , so that a WZ term appears in $S_g(i)$ for all i . The effective action S_{JJA} for the array is found by adding together the action for each tunnel junction $S_{JJ}(i, n_i)$;

$$S_{JJA} = \sum_i \left\{ S_0(i) + S_{wz}(i) + S_2(i) + \sum_{\{n_i\}} [S_c(i, n_i) + S_t(i, n_i)] \right\} \quad . \quad (4)$$

The current drive contribution to $S_t(i, n_i)$ is obtained from eqn. (3) by the substitution $\gamma \rightarrow \gamma(i, n_i)$; $\hat{\mathbf{n}} \rightarrow \hat{\mathbf{n}}(i, n_i)$. Note that $\dot{\mathbf{r}}_0$ will not be perpendicular to *all* $\hat{\mathbf{n}}(i, n_i)$ so that the Magnus drive *is* active in a JJA. We now consider the Continuum Limit (CL) for the SC dynamics of the JJA. Let l be the length scale over which the gauge invariant phase difference $\gamma(i, n_i)$ varies; the CL of the SC dynamics corresponds to $l \gg a_0$. In this limit $\gamma(i, n_i)$ varies little from grain to grain. For an array in the CL, it is reasonable to coarse grain S_{JJA} over a length scale d satisfying $a_0 \ll d \ll l$. Let the JJA map onto the region $A \subset R^2$, and partition A into cells $dA(\mathbf{x})$ of area d^2 . In the CL, $\gamma(i, n_i)$ is essentially constant over each cell. All grains and junctions lying within $dA(\mathbf{x})$ will be grouped together in eqn. (4) to give the contribution from this cell to the coarse grained JJA action $\bar{S}_{JJA}$. Because of the JG-eqn, $S_c(i, n_i)$ is seen to represent the action associated with the electric field $\mathbf{E}(i, n_i)$ present in this junction. A similar term is present in $S_2(i)$ (see eqn. (1)) so that the sum of these two types of terms, for the cell at $\mathbf{x}$, gives the action associated with the electric field present in this cell. The coarse graining of these terms gives the action $dA(\mathbf{x})[-\bar{\mathbf{E}}^2/8\pi]$ associated with the coarse grained electric field $\bar{\mathbf{E}}(\mathbf{x})$ in this cell. One can show [6] that, in the CL, the integrand of $S_t(i, n_i)$ is proportional to $\mathbf{v}_s^2(i, n_i)$. A similar term is present in $S_2(i)$ so that the sum of these two types of terms for the cell at

$\mathbf{x}$ gives the action associated with the superflow kinetic energy in this cell. As with the electric field, coarse graining leads to the action $dA(\mathbf{x})[m\bar{\rho}_s(\mathbf{x})\bar{\mathbf{v}}_s^2(\mathbf{x})/2]$ associated with the coarse grained superflow $\bar{\mathbf{v}}_s(\mathbf{x})$. Coarse graining $S_0(i)$, $S_{wz}(i)$, and the remaining terms in $S_2(i)$ is straightforward as they are essentially constant over a cell $dA(\mathbf{x})$ and so yield terms of similar form, but with coarse grained quantities appearing in the final result. We find,

$$S_{JJA} \rightarrow \bar{S}_{JJA} = S_0 + \int_A d^2x dt \left[\bar{\rho}_s \left(\frac{\hbar}{2} \partial_t \phi + e\bar{A}_0 \right) + N(0) \left(\frac{\hbar}{2} \partial_t \phi + e\bar{A}_0 \right)^2 + \frac{m\bar{\rho}_s}{2} \bar{\mathbf{v}}_s^2 + \frac{1}{8\pi} \left\{ (\bar{\mathbf{H}} - \mathbf{H}_{ext})^2 - \bar{\mathbf{E}}^2 \right\} \right].$$

Thus $\bar{S}_{JJA}$ has exactly the same form as the action of a single bulk 2D superconductor S_g (see eqn. (1)), so that, in the CL, the SC dynamics of a JJA maps onto that of a 2D SC film. Because the Berry phases are non-trivial in the array, $\bar{S}_{JJA}$ will contain a WZ term, and consequently, the non-dissipative force $\mathbf{F}_{nd}$ acting on a vortex will contain a Magnus force contribution, exactly as in the 2D film [1,4]. We stress that the appearance of the Magnus force in the dynamics of vortices in a JJA follows naturally from the presence of a Berry phase in the SC dynamics; it is not necessary to assume that external charges are fixed to the grains to generate the Magnus force as done in Ref. [12]. Having established that the vortex dynamics of a JJA maps onto that of a 2D SC film, we must point out an important advantage of arrays in the CL over 2D films for studies of vortex dynamics. Because the array lattice can be constructed with very great precision, one can control the degree of disorder present in a particular array. This raises the possibility of carrying out flux-flow experiments in arrays (in the CL) in which the degree of disorder can be systematically varied from (essentially) zero to any specified degree, or distribution, of disorder. This is usually *not* the case in 2D films. As an example, consider the Hall Effect in an array in the CL with negligible disorder. The only forces acting on a vortex will be the non-dissipative force $\mathbf{F}_{nd} = a\mathbf{v}_s \times \hat{\mathbf{z}} - b\mathbf{v}_L \times \hat{\mathbf{z}}$ (a, b being determined via the coarse grain averaging); and the dissipative force $-\eta\mathbf{v}_L$. In the steady state, the sum of these forces is zero, allowing a determination of the Hall angle, $\tan \Theta_H = -v_{Lx}/v_{Ly} = b/\eta$. This is exactly the result one finds in a 2D SC film [7]. Thus one expects a sign anomaly in the Hall Effect to occur in JJA's (in the CL) when the grains are made of a material which shows a sign anomaly in the 2D film experiments. The ability to systematically vary disorder in arrays would appear to make them ideally suited for testing two recent predictions of Vinokur et. al. [7]: (i) independence of the Hall conductivity σ_{xy} on disorder; and (ii) power law behavior of the Hall resistivity $\rho_{xy} \sim \rho_{xx}^2$ after averaging over disorder.

We would like to thank the International Atomic Energy Agency; UNESCO; and ICTP, Trieste for support. F. G. would like to thank T. Howell III for constant support.

-
- [1] P. Ao and D. J. Thouless, Phys. Rev. Lett. **70**, 2158 (1993).
 - [2] Regrettably, what is meant by the ‘‘Magnus Force’’ is not uniformly agreed upon in the literature of Type-II superconductors. In this paper we will follow the terminology introduced in Ref. [4].
 - [3] M. Stone, *Bosonization* (World Scientific, New Jersey, 1994), chap. 3; I. J. R. Aitchison, P. Ao, D. J. Thouless, and X.-M. Zhu, Phys. Rev. B **51**, 6531 (1995).
 - [4] F. Gaitan, Phys. Rev. B **51**, 9061 (1995).
 - [5] Thermodynamic manifestation of Berry’s phase in the magnetization of superconducting networks in the mixed state has been discussed in R. Rammal, Physica B **152**, 37 (1988).
 - [6] F. Gaitan and S. R. Shenoy, in preparation.
 - [7] V. M. Vinokur, V. B. Geshkenbein, M. V. Feigel’man, and G. Blatter, Phys. Rev. Lett. **71**, 1242 (1993).
 - [8] U. Eckern, G. Schon, and V. Ambegaokar, Phys. Rev. B **30**, 6419 (1984).
 - [9] P. W. Anderson, Rev. Mod. Phys. **38**, 298 (1966).
 - [10] B. D. Josephson, Phys. Lett. **1**, 251 (1962). Damping due to quasiparticle tunneling is also introduced by the tunneling interaction; it will be ignored as it does not affect our arguments.
 - [11] B. Dueholm et. al. , in *Proceedings of the Seventeenth International Conference on Low Temperature Physics* , eds. U. Eckern, A. Schmid, W. Weber, and H. Wühl (North Holland, Amsterdam 1984), p. 691.
 - [12] R. Fazio et. al. , Helv. Phys. Acta **65**, 228 (1992).